
Joint first authors

Cochrane permits adding joint first authors for when two or more authors may agreed to have contributed equally to a review and wish to share the position of first author in the byline.

This is displayed in the *Cochrane Database of Systematic Reviews* in the author information; such as shown in https://doi.org/10.1002/14651858.CD009593.pub4:

HTML display



PDF display



Xpert MTB/RIF and Xpert MTB/RIF Ultra for pulmonary tuberculosis and rifampicin resistance in adults

David J Horne[1a], Mikashmi Kohli[2b], Jerry S Zifodya[3], Ian Schiller[4], Nandini Dendukuri[4], Deanna Tollefson[5], Samuel G Schumacher[6], Eleanor A Ochodo[7], Madhukar Pai[2], Karen R Steingart[8]

[1]Department of Medicine, Division of Pulmonary and Critical Care Medicine, and Firland Northwest TB Center, University of Washington, Seattle, WA, USA. [2]Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Canada. [3]Pulmonary and Critical Care Medicine, University of Washington, Seattle, WA, USA. [4]Division of Clinical Epidemiology, McGill University Health Centre - Research Institute, Montreal, Canada. [5]Department of Global Health, University of Washington, Seattle, WA, USA. [6]FIND, Geneva, Switzerland. [7]Centre for Evidence-based Health Care, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa. [8]Honorary Research Fellow, Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK

[a]These authors contributed equally to this work. [b]These authors contributed equally to this work

For Editors: see Joint first authors: assigning in RevMan